Research and Monitoring: Evolution, Methods and Case Studies



Jeff Miller

William J Miller@nps.gov

Matt Patterson

Dr. Andrea Atkinson Judd Patterson Rob Waara Dr. Kevin R.T. Whelan Brian Witcher Alexandra Wright

Lessons learned from seventeen years of monitoring coral reefs



Lessons learned

1. Worthwhile endeavor


2. Takes time and commitment

3. Be flexible to react to change driving events










































































"It requires effort to ensure that sampling is why bother?" is random."

- Roger Green
Sampling Design and Statistical Methods
for Environmental Biologists



Potential problems with haphazard sampling designs

- Poor Inference: Use of judgment to select sites, convenience sampling
 results only apply to pre-selected sites
- Poor Stratification: stratification is based upon strata that change through time. Includes placement of sites in temporary hot spots
 -> analysis complication, validity of trends open to question
- Design does not match objectives: Issues of sufficient power, appropriate scope, and appropriate link between protocol and indicator
 - -> unable to answer managers' real questions in time frame needed
 - -> or needless wasting of effort on detailed answers for a ill-defined question
- Not Robust: changes in personnel or protocol or funding affect interpreting data
 - -> analysis complication, validity of trends open to question



Lessons learned

1. Worthwhile endeavor

2. Takes time and commitment

3. Be flexible to react to change driving events





Pilot program phase (3-5 years)


- Piloting protocols
 - Establishing variability, evaluating power & optimizing design & protocol
- Revising protocol & sampling designs
- Establishing baselines
- Long-term monitoring to evaluate change




Virgin Islands National Park - STJ
























Buck Island Reef National Monument











South Florida National Parks

Biscayne NP




Amanda's Reef Biscayne NP Sampled: 2004-2006 Area= 20,240 m²



Ball Buoy Reef
Biscayne NP
Sampled: 2004-2006
Area= 14,136 m²

Dry Tortugas NP

















Some important questions for scientists & management...

- What amount of change do you want to be able to detect?
- How confident do you need to be that you'll catch this change the first sampling event it occurs? (90%? 70%?)
- What level of "false alarms" are you willing to except? (5%? 10%?)




Power – detect trend 25% over 5 years


 $(\alpha = 0.05, 20 \text{ permanent transects})$


		Mean
Site (Park/Location)	Power	Coral Cover
		40.007
Newfound Reef (STJ)	92%	13.6%
Yawzi Reef (VIIS)	74%	7.3%
Haulover Reef (VIIS)	96%	22.3%
Mennebeck Reef (VIIS)	97%	23.7%
W. Spur and Groove (BUIS)	76%	5.2%
S. Fore Reef (BUIS)	99%	17.9%
Amanda's Reef (BISC)	57%	7.0%
Ball Buoy Reef (BISC)	70%	6.6%
Bird Key Reef (DRTO)	91%	12.9%



Found power and recommended sample sizes varied with average coral cover











Sample Size Comparison Permanent vs. Non-permanent Transects

 $(\alpha = 0.05)$





20 non-permanent transects: 21% power

92 non-permanent transects: 80% power

20 non-permanent transects: 14% power

181 non-permanent transects: 80% power



Lessons learned

1. Worthwhile endeavor

2. Takes time and commitment

3. Be flexible to react to change driving events



2005 Episadie Manttonigg Event

S. Fore Reef, BUIS


Tektite, VIIS

Haulover, VIIS

Mennebeck, VIIS

Yawzi, VIIS

Newfound, STJ



95.6% coral cover bleached

90.2% coral cover **90.2%** coral


96.4% coral cove blooms

93.6% coral cover bleached

Bleached cover bleached

91.8% coral cover bleached

Establishing baseline allows testing for both sudden & gradual changes














Successful and vital partnership with USGS


Began 2005 co-locating benthic cover with mortality from disease on permanent transects

Co-location of benthic cover and mortality from disease











Bleaching – Disease connection








August

September

October







November

December

January




23% loss

In one

Year!

48% loss 24% loss 28% loss

Tektite - in six months!!



Stakel Solde häßuy-in"








Photo by Judd Patterson




Bleaching – Disease connection



2005



Disease




2006





Coral Bleaching Recovery - individual coral tracking






























DRTO Bird Key Reef

Sampled: 2004 (20 ten-meter random transects)


Area = $26,365 \text{ m}^2$








DRTO Bird Key Reef Sampled: 1997 and 2004 Haphazard vs. Random sampling












DRTO Bird Key Reef Sampled: 1997 and 2004

Repeated Sampling of transect between permanent pins











DRTO Bird Key Reef Sampled: 1997 and 2004

Repeated Sampling of transect between permanent pins







Summary

- Permanent visual record
- Quantitative and Qualitative data
- •Electronic transfer of data and images
- •Repeatable, "Power"-ful, rigorous
- •Relatively simple and efficient
- •Results and images for scientific community and policy-makers







Thank you for your attention.

Any questions?

